Math 138 – Practice midterm

Due date: October 15, 2025

Instructions:

- You have 2 hours to complete this exam.
- No external resources are allowed.
- Upload your completed practice midterm to Crowdmark.

Word of advice: While not 100%, we can likely tell if you use AI on this exam. Ultimately the grade on the practice midterm is *completion based*. To cheat on this will likely not result in a higher score on the practice midterm, but will almost certainly result in a lower score on the actual exam. Similarly, the two hour limit is not enforceable, but you should still follow it as it will help you get a better sense of your preparedness for the actual exam.

Name: _____

Question 1. (15 pts)

Spot the error in the following proof, and explain why it is wrong.

Proposition. Every function $f: \{1, ..., n\} \rightarrow \{1, ..., n\}$ has a fixed point (i.e., for some x one has f(x) = x).

Proof. We prove this by induction on n. When n=1 the only function $f:\{1\} \to \{1\}$ satisfies f(1)=1, so has a fixed point. For the induction hypothesis, assume the claim is true for any function $g:\{1,\ldots,n\} \to \{1,\ldots,n\}$, and consider a function

$$f: \{1, \dots, n+1\} \to \{1, \dots, n+1\}.$$

Case 1: If f(n+1) = n+1 we're done.

Case 2: If $f(n+1) \neq n+1$, we can consider the function $g: \{1, \ldots, n\} \to \{1, \ldots, n\}$ given by f(i) = g(i). By induction hypothesis there is some $x \in \{1, \ldots, n\}$ such that x = g(x) = f(x).

Thus, in either case we see f has a fixed point as desired.

Question 2. (15 pts)

Let $n \geqslant k \geqslant 0$ be integers. Prove that

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

Question 3. (20 pts)

Suppose that $x \in \mathbb{Q}$. Use the *p*-adic valuation to prove that if $x^{\frac{2}{3}}$ belongs to \mathbb{Q} then $x^{\frac{1}{3}}$ belongs to \mathbb{Q} .

Question 4. 20 pts

Let A and B be subsets of a set S. Prove that

$$A\triangle B = (A \cup B) \cap (A^c \cup B^c).$$

Math	138 -	Practice	midterm
maun	100	1 lactice	muaterm

Question 5. (30 pts)

Prove by induction that if T_n is the number of length $n \ge 1$ strings in the symbols 0, 1, 2 with exactly one 2 then

$$T_n = n \cdot 2^{n-1}$$